

1. (11 points)

5

- (a) The number of H1N1 viruses H (measured in trillions) that infect a college student is a function of the degree of immunosuppression I (the fraction of the immune system that is turned off by studying) according to $H(I) = 6I^2 + 2$. The fever F (measured in $^{\circ}\text{C}$) associated with an infection is a function of the number of viruses according to $F(H) = 37 + 0.5H$. Find fever as a function of immunosuppression. What will the fever be if immunosuppression is complete ($I = 1$)?

$$H(I) = 6I^2 + 2$$

$$F(H) = 37 + 0.5H$$

Find $F(I)$:

$$\begin{aligned} F(H(I)) &= F(6I^2 + 2) = 37 + 0.5(6I^2 + 2) = 37 + 3I^2 + 1 \\ &= 38 + 3I^2; \end{aligned}$$

If $I = 1$ $F(1) = 38 + 3 = 41$

$$F(I) = 38 + 3I^2$$

- (b) Consider the data in the following table describing the number of students in Dorm Q that come down with the H1N1 flu as a function of the average number of hours that students in the dorm spend studying each day.

x	average number of hours per day spent studying	number of students sick with H1N1	y
	1	34	
	4.5	13	
	5	10	

4

- i. These data lie on a line (to convince yourself of this, you may graph the data, but you do not need to show a graph here). Find the equation of the line connecting the first two points.

$$y - y_0 = m(x - x_0). \quad (x_0, y_0) = (1, 34)$$

$$m = \frac{13 - 34}{4.5 - 1} = \frac{-21}{3.5} = -6$$

equation: $y - 34 = -6(x - 1); \quad \underline{y = -6x + 40}$

2

- ii. How many students does your equation predict would get sick in in Dorm Q if the average numbers of hours that students in this dorm spend studying each day is 8? Does this make sense? Why or why not?

$$y = -6(8) + 40 = -8 < 0$$

No, this does NOT make sense since students don't come in negatives.

2. (12 points) Consider the discrete-time dynamical system with updating function

$$b_{t+1} = 0.4b_t$$

representing the population of bacteria.

5

(a) Find the solution of the discrete-time dynamical system if $b_0 = 1.0 \times 10^4$.

$$b_t = 1.0 \times 10^4 (0.4)^t \\ (= 1.0 \times 10^4 e^{\ln(0.4)t})$$

7

(b) Find the half-life of the population.

$$\frac{1}{2} (\cancel{1.0 \times 10^4}) = \cancel{1.0 \times 10^4} (0.4)^t \\ \frac{1}{2} = 0.4^t \\ \ln\left(\frac{1}{2}\right) = \ln(0.4^t) \\ \ln\left(\frac{1}{2}\right) = t \ln(0.4); \\ t = \frac{\ln\left(\frac{1}{2}\right)}{\ln(0.4)} \\ \approx \underline{\underline{0.756}}$$

or:

$$\frac{1}{2} \cancel{b_0} = \cancel{b_0} e^{\ln(0.4)t} \\ \frac{1}{2} = e^{\ln(0.4)t} \\ \ln\left(\frac{1}{2}\right) = \ln(0.4)t; \\ t = \frac{\ln\left(\frac{1}{2}\right)}{\ln(0.4)}$$

3. (14 points) Suppose that 40 percent of the medication in a patient's body is **removed** each day by the body but the patient takes an additional dose of 2 mg/L at the end of each day.

- (a) Write the updating function $M_{t+1} = f(M_t)$ for the concentration of medicine on day $t+1$ as a function of the concentration on day t . Recall that

$$\text{new concentration} = \text{old concentration} - \text{fraction used} \times \text{old concentration} + \text{supplement.}$$

4

$$M_{t+1} = M_t - 0.4M_t + 2 = 0.6M_t + 2$$

$$M_{t+1} = 0.6M_t + 2$$

- (b) Find the equilibrium point algebraically, if it exists.


4

$$M^* = 0.6M^* + 2$$

$$0.4M^* = 2$$

$$M^* = \frac{2}{0.4} = 5$$

- (c) Draw the graph of the updating function, identify the equilibrium if it exists, and cobweb the concentration for days 1, 2, and 3 if the initial dose is $M_0 = 10$ mg/L. (Make sure to label clearly all points (M_t, M_{t+1}) on the updating function during cobwebbing.)



- 2
- (d) If $M_0 = 10$, the solution to this discrete time dynamical system is $M_t = 5 \cdot (1 + (\frac{3}{5})^t)$ where t is time, measured in days. While the patient requires a high initial dose the patient will be at risk if the concentration remains above 6 mg/L for 3 days or more. Use the solution given to determine if this patient will be at risk. (Hint: What is M_3 ?)

$$M_3 = 5(1 + (\frac{3}{5})^3) = 6.08 > 3.$$

yes, The patient is indeed at risk.

4. (12 points)

(a) Find all non-negative equilibria of the discrete-time dynamical system

$$m_{t+1} = \frac{2m_t}{1 + am_t},$$

where a is a positive parameter. How many equilibria are there if $a = 0$?

$a > 0:$ $m^* \doteq \frac{2m^*}{1 + am^*}$

$$m^*(1 + am^*) = 2m^*$$

$$m^* + a(m^*)^2 = 2m^*$$

$$a(m^*)^2 - m^* = 0$$

$$m^*(am^* - 1) = 0$$

$$m^* = 0 \text{ or}$$

$$am^* - 1 = 0; m^* = \frac{1}{a}.$$

equilibria:

$m^* = 0 \text{ and}$
 $m^* = \frac{1}{a}$


If $a = 0,$

$$m^* = \frac{2m^*}{1};$$

$m^* = 0$ is the only equilibrium.

1

(b) The updating function for a discrete-time dynamical system $x_{t+1} = f(x_t)$ is graphed below. Use cobwebbing to help you label each of the three equilibria as stable or unstable.




5. (12 points) Let V_{t+1} represent the voltage of the AV node in the heart model.


$$V_{t+1} = \begin{cases} e^{-\alpha\tau}V_t + u, & \text{if } V_t \leq e^{\alpha\tau}V_c \\ e^{-\alpha\tau}V_t, & \text{if } V_t > e^{\alpha\tau}V_c \end{cases}$$

(a) For each of the following two graphs of the updating function, cobweb starting from an initial value of $V_0 = 10$ and determine if the heart is healthy, has 2:1 AV block, or the Wenckebach phenomenon.

4



Healthy



more beats than skips;

Wenckebach phenomenon

(b) Let $e^{-\alpha\tau} = 0.6$, $V_c = 21$, and $u = 8$. If $V_0 = 26$ will the heart beat? Why or why not? Calculate V_1 .

4

$$e^{\alpha\tau}V_c = \frac{21}{0.6} = 35.$$

Since $V_0 = 26 < 35$, yes the heart will beat.

$$V_1 = 0.6(26) + 8 = \boxed{23.6 = V_1}$$

(c) Does the system described in part b have an equilibrium? Why or why not? If it has an equilibrium, find it algebraically.

4

If there is an equilibrium V^* then

$$V^* = e^{-\alpha\tau}V^* + u = 0.6V^* + 8;$$

$$0.4V^* = 8;$$

$$\underline{V^* = 20}$$

Since $V^* = 20 < e^{\alpha\tau}V_c$

$= \frac{21}{0.6}$, V^* is indeed an equilibrium; yes, the system has an equilibrium.


6. (13 points) (a) Find the following limits, if they exist. Show all of your work and justify your answers to receive full credit. If a limit does not exist, write "DNE," and explain why it does not exist.

2 i) $\lim_{x \rightarrow 1} \frac{7x^2 + 3x}{2x - 1} = \frac{7(1)^2 + 3(1)}{2(1) - 1} = \underline{10}$

2 ii) $\lim_{t \rightarrow -2} \frac{t^2 + 9t + 14}{t + 2} = \lim_{t \rightarrow -2} \frac{\cancel{(t+2)}(t+7)}{\cancel{t+2}} = \lim_{t \rightarrow -2} (t+7) = \underline{5}$.

2 iii) $\lim_{z \rightarrow 2} \frac{z}{(z-2)(z+2)}$

Graph:




DNE since

$$\lim_{z \rightarrow 2^+} \frac{z}{(z-2)(z+2)} = +\infty$$

$$\lim_{z \rightarrow 2^-} \frac{z}{(z-2)(z+2)} = -\infty$$


2 iv) $\lim_{y \rightarrow 0} \frac{1}{y^4} = \infty$.



$$\lim_{y \rightarrow 0^+} \frac{1}{y^4} = \infty = \lim_{y \rightarrow 0^-} \frac{1}{y^4}$$

(continuation of problem 6)

(b) For the function $f(x)$ graphed below, find the following limits or explain why the limit does not exist. (If a limit exists, you may just give a number without explanation.)



1 point
apiece

a) $\lim_{x \rightarrow 3} f(x) = 5$

b) $\lim_{x \rightarrow 4^-} f(x) = 6$

c) $\lim_{x \rightarrow 4^+} f(x) = 3$


d) $\lim_{x \rightarrow 4} f(x)$ DNE since $\lim_{x \rightarrow 4^+} f(x) = 3 \neq 6 = \lim_{x \rightarrow 4^-} f(x)$

e) $\lim_{x \rightarrow 6} f(x) = 8$

7. (12 points) If you use a calculator, please make sure that it is in radians.

(a) Accurately graph the function

$$M(t) = \begin{cases} \cos\left(\frac{\pi}{2}t\right), & t < 4 \\ 0.5t - 1, & t \geq 4 \end{cases}$$



(b) Does $\lim_{t \rightarrow 4} M(t)$ exist? If so, find the value of the limit. If not, explain why not.

4

$$\lim_{t \rightarrow 4^-} M(t) = \lim_{t \rightarrow 4^-} \cos\left(\frac{\pi}{2}t\right) = \cos(2\pi) = 1.$$

$$\lim_{t \rightarrow 4^+} M(t) = \lim_{t \rightarrow 4^+} (0.5t - 1) = 0.5(4) - 1 = 1.$$

$$\therefore \boxed{\lim_{t \rightarrow 4} M(t) = 1}$$

(c) Is $M(t)$ continuous at $t = 4$? If so, use the definition of continuity to explain how you know this. If not, explain why it is not continuous there.

4
Yes, since

(i) $M(4)$ is defined ($M(4) = 1$).

(ii) $\lim_{t \rightarrow 4} M(t)$ exists ($\lim_{t \rightarrow 4} M(t) = 1$)

∴ (iii) $\lim_{t \rightarrow 4} M(t) = M(4)$.

8. (14 points) Tom Brady forgot to shower after a football game, so foot fungus began to grow. Suppose that the fungus population $f(t)$ on Tom's toes at time t is given by $f(t) = 4t^2 - 4t + 2$.

- (a) Find a formula for the slope of the secant line that passes through $(2, f(2))$ and $(2 + \Delta t, f(2 + \Delta t))$.

4

$$\begin{aligned} \text{slope} &= \frac{f(2 + \Delta t) - f(2)}{(2 + \Delta t) - 2} = \frac{[4(2 + \Delta t)^2 - 4(2 + \Delta t) + 2] - [4(2)^2 - 4(2) + 2]}{\Delta t} \\ &= \dots = \frac{12\Delta t + 4\Delta t^2}{\Delta t} = 12 + 4\Delta t. \end{aligned}$$

- (b) Find the average rate of change in the fungus population between times $t = 2$ and $t = 2.1$.

3

Average Rate of Change = Slope of Secant Line.
 Using the result from part a, with $\Delta t = (2.1 - 2) = 0.1$,
 average rate of change between $t = 2$ and $t = 2.1$
 $= 12 + 4(0.1) = \underline{12.4}$

- (c) Use the definition of the derivative to find $f'(2)$.

3

$$f'(2) = \lim_{\Delta t \rightarrow 0} \frac{f(2 + \Delta t) - f(2)}{(2 + \Delta t) - 2} \stackrel{\text{part a}}{=} \lim_{\Delta t \rightarrow 0} (12 + 4\Delta t) = \underline{12}.$$

- (d) What is the equation of the tangent line to $f(t)$ at time $t = 2$?

4

The tangent line is given by

$$y - f(2) = f'(2)(x - 2);$$

$$y - 10 = 12(x - 2);$$

$$y = 12x - 24 + 10;$$

$$\boxed{y = 12x - 14}$$